### Directions for use Read carefully!



# **ECOSOL G5** (Solution for **Intravenous Infusion**)

#### Symptoms and treatment for overdosage

If hyperglycaemia or glucosuria occur, the dosage and infusion rate should be reduced, or an insulin administration is recommended.

#### Side effects

Hyperglycemia and renal loss may occur in case of reduced glucose tolerance. These manifestations are normally prevented by reducing the dosage and/or by giving insulin. Enhanced bilirubin and lactate levels may be found if the recommended dosage is exceeded. Glucose solutions are acidic (pH 3.5 - 5.0) and may produce thrombophlebitis.

#### Contraindications

Diabetes mellitus (with the exception of hypoglycaemia conditions) Glucose intolerance Hypotonic dehydration if lacking electrolytes are not replaced Overhydration Hypokalaemia

#### Drug interaction

The compatibility of any additives to this solution should be checked before use.

> Manufactured by: B. Braun Medical

Imported by: PT. B. Braun Medical

Indonesia

Industries Sdn. Bhd. 11900 Bayan Lepas, Penang, Malaysia. Licence Holder:

Storage Do not store above 30°C.

Presentation Plastic bottle 100 ml, 500 ml

Reg no.: DKI1277201749A1

Harus dengan resep dokter

#### schwarz

 $Dokument = 148 \times 210 mm$ DIN A 5 - 1 Page

ID 4 GIF - EP (Unfolded) 4/12626137/0718 Production site: Penang

Lätus: 3031



Font size: 8 pt.

G 160676

#### Composition

Each 100 ml solution contains:

Glucose Monohydrate for Parenteral Use Water for Injections to		5.5 g 100 ml
Caloric Value	kJ/I	850
	kcal/l	200
Osmolarity	mOsm/l	278

#### Mechanism of action

Glucose is absorbed from the gastrointestinal tract, and oxidized as a source of energy, or stored in the liver as glycogen. It is the only energy substrate which is directly, instantly and universally utilized by the body. Glucose is vital for the myocardium, brain and nerves.

#### Indications

Energy supply Hypertonic dehydration Vehicle solution for supplementary medication

#### Posology

The maximum dosage is 500 mg/kg/hr (10 ml/kg/hr of Glucose 5 %) over period of less than 24 hours.

#### Route of administration: I.V.

#### Precautions/Warning

Blood glucose, serum electrolytes and water balance should be monitored regularly.

Electrolytes are to be supplemented as required. The compatibility of any additives to these solutions should be checked before use. Glucose solution should not be administered through the same infusion set through which also blood has or may be given because of the risk pseudoagglutination.

#### Cautions

The compatibility of any additives to this solution should be checked before use.

## **B BRAUN**



