

**NAME OF THE MEDICINAL PRODUCT**

Compound Sodium Lactate Solution for Infusion  
(Hartmann's Solution)

**COMPOSITION**

1000 ml of solution for infusion contain:

Sodium chloride	6.00 g
Sodium lactate solution (50% w/w) (equivalent to sodium lactate, 3.12 g)	6.24 g
Potassium chloride	0.40 g
Calcium chloride dihydrate	0.27 g

*Electrolyte concentrations:*

Sodium	131 mmol/l
Potassium	5.4 mmol/l
Calcium	1.8 mmol/l
Chloride	112 mmol/l
Lactate	28 mmol/l

*Excipient:* Water for injection.

**THERAPEUTIC INDICATIONS**

Fluid substitution under the conditions of undisturbed acid-base balance or mild acidosis, isotonic and hypotonic dehydration, short-term intravascular volume replacement, vehicle solution for compatible electrolyte concentrates and drugs.

**CONTRAINDICATIONS**

Impairment of lactate utilisation with hyperlactatemia, hyperhydration.

**UNDESIRABLE EFFECTS**

Undesirable effects are listed according to their frequencies as follows:

Not known: (cannot be estimated from the available data)

***Metabolism and nutrition disorders***

Not known: Hospital Acquired Hyponatremia

***Neurological disorders:***

Not known: Hyponatremic encephalopathy

Hospital acquired hyponatremia may cause irreversible brain injury and death due to development of acute hyponatremic encephalopathy.

**WARNINGS**

Keep out of sight and reach of children. For single use only. Discard container and unused content after use. Use only if solution is clear, colourless and container and closure are undamaged. Expel all air before starting pressure infusion.

**NOTE**

Not all products are registered and approved for sale in all countries or regions. Indications of use may also vary by country and region. Please contact your country representative for product availability and information.

**B. Braun Melsungen AG, 34209 Melsungen, Germany, 01/2018**