

## **NAME OF THE MEDICINAL PRODUCT**

Glucose B. Braun 50 mg/ml solution for Infusion

## **COMPOSITION**

1 ml of solution contains 50 mg glucose (as glucose monohydrate 55.0 mg)  
100 ml of solution contain Glucose 5.0 g (as glucose monohydrate, 5.5 g)

*Excipient:* Water for injection.

## **THERAPEUTIC INDICATIONS**

Carbohydrate solution for intravenous liquid therapy; vehicle solution for compatible medicinal products.

## **CONTRAINDICATIONS**

Hypersensitivity to the active substance, hyperglycemia, not responding to insulin doses of up to 6 units insulin/hour, lactic acidosis.

If it is necessary to administer large volumes, further contraindications can arise on account of the fluid load: Hypotonic hyperhydration, isotonic hyperhydration, acute congestive heart failure, pulmonary edema.

This solution must not be used alone for fluid supply/rehydration because it does not contain electrolytes.

## **UNDESIRABLE EFFECTS**

Undesirable effects are listed according to their frequencies as follows:

Not known: (cannot be estimated from the available data)

### ***Immune system disorders***

Not known: Anaphylactic reaction\*, hypersensitivity\*

***Skin and subcutaneous tissue disorders***

Not known: Rash

***Vascular disorders***

Not known: Venous thrombosis, phlebitis

***General disorders and administration site conditions***

Not known: Chills\*, pyrexia\*, infusion site infection, infusion site irritation for example erythema, extravasation, local reaction, pain localised

***Metabolism and nutrition disorders***

Not known: Electrolyte imbalance, e.g. hyponatremia and hypokalemia, hospital acquired hyponatremia\*\*

***Neurological disorders***

Not known: Hyponatremic encephalopathy\*\*

\*Potential manifestation in patients with corn allergy

\*\* Hospital acquired hyponatremia may cause irreversible brain injury and death due to development of acute hyponatremic encephalopathy

***Reporting of suspected adverse reactions***

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system.

**WARNINGS**

Keep out of the sight and reach of children. For single use only. Discard unused contents.  
Do not reconnect partially used containers. Use only if solution is clear and colourless or almost colourless and the container and closure are undamaged.

**NOTE**

Not all products are registered and approved for sale in all countries or regions. Indications of use may also vary by country and region. Please contact your country representative for product availability and information.

**B. Braun Melsungen AG, 34212 Melsungen, Germany, 06/2018**