

## NAME OF THE MEDICINAL PRODUCT

Sodium Chloride 5.85 % Braun Concentrate for Solution for Infusion

## COMPOSITION

1 ml of concentrate for solution for infusion contains

Sodium chloride 58.5 mg

*Electrolytes:*

Sodium 1 mol/l

Chloride 1 mol/l

1 ampoule of 20 ml contains 1.17 g Sodium chloride

1 vial of 100 ml contains 5.85 g Sodium chloride

*Excipient:* Water for injection.

## THERAPEUTIC INDICATIONS

Hyponatremia; hypochloremia; hypotonic hyperhydration.

## CONTRAINDICATIONS

Hypernatremia, hyperchloremia.

## UNDESIRABLE EFFECTS

The frequency of possible adverse reactions primarily depends on the dosage and the infusion rate.

***Metabolism and nutrition disorders***

Hypernatremia and hyperchloremia may occur during application.

Excessively rapid infusion of solutions with a high sodium chloride concentration may lead to acute volume overload.

***Gastrointestinal disorders***

Excessively rapid infusion of solutions with a high sodium chloride concentration may lead to diarrhoea.

***Renal and urinary disorders***

Excessively rapid infusion of solutions with a high sodium chloride concentration may lead to osmotic diuresis.

***Administration site conditions***

Peripheral vein administration of solutions with a high sodium chloride concentration may lead to irritation and inflammation of the vein septum.

**WARNINGS**

Keep out of the sight and reach of children. Sterile and free of bacterial endotoxins. Intravenous application after dilution. For single use only. Discard unused contents. Only to be used if the solution is clear and container and closure are undamaged. Do not store above 25 °C.

**NOTE**

Not all products are registered and approved for sale in all countries or regions. Indications of use may also vary by country and region. Please contact your country representative for product availability and information.

**B. Braun Melsungen AG, 34212 Melsungen, Germany, 07/2014**